

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**




B5

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 37/02, 39/00, 39/12		A1	(11) International Publication Number: WO 92/15318
			(43) International Publication Date: 17 September 1992 (17.09.92)
(21) International Application Number: PCT/US92/01705 (22) International Filing Date: 5 March 1992 (05.03.92) (30) Priority data: 665,762 7 March 1991 (07.03.91) US (71) Applicant: SERAGEN, INC. [US/US]; 97 South Street, Hopkinton, MA 01748 (US). (72) Inventor: NICHOLS, Jean, C. ; 15 Astra Way, Wayland, MA 01778 (US). (74) Agent: CLARK, Paul, T.; Fish & Richardson, 225 Franklin Street, Boston, MA 02110-2804 (US).		(81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL (European patent), NO, PL, RO, RU, SD, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent). Published With international search report.	

(54) Title: USE OF CELL SURFACE RECEPTOR TARGETED MOLECULES FOR THE TREATMENT OF VIRAL DISEASES



(57) Abstract

The invention features a method for treating a patient infected with a virus. The method includes administering to the patient a molecule which is capable of specifically binding to a proteinaceous cell receptor expressed on a cell of the patient which cell contributes to the disease state of the patient, the molecule being capable of decreasing the viability of the cell.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LJ	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

- 1 -

**USE OF CELL SURFACE RECEPTOR TARGETED MOLECULES
FOR THE TREATMENT OF VIRAL DISEASES**

Background of the Invention

5 This invention relates to the treatment of diseases associated with viral infection.

 Human immunodeficiency virus (HIV), the etiologic agent of acquired immunodeficiency syndrome (AIDS), is a retrovirus which selectively infects certain immune
10 system cells, including T4 (CD4+) lymphocytes and CD4+ cells of the monocyte/macrophage lineage. In its advanced stages, HIV infection causes immune system failure and renders the victim susceptible to opportunistic infections and neoplasms. In the absence
15 of effective treatment, the mortality rate for AIDS patients approaches 100% (Fauci, *Science* 239:617, 1988).

 T4 lymphocytes play a central role in many immune system functions, and the cytopathic effect of HIV on infected T4 lymphocytes is thought to be responsible for
20 the devastating effect of HIV infection. Active viral replication usually leads to the death of the host cell. However, in certain host cells the virus does not immediately replicate, and these cells become chronically or latently infected (Fauci, *supra*). Chronically
25 infected cells express viral proteins at a low level; latently infected cells have an integrated provirus but do not express viral proteins. Although viral replication does not occur in chronically or latently infected cells until the cells are activated, these cells
30 can serve as a reservoir of HIV in the body.

 Previous approaches to slowing the progression of HIV infection include the use of antibodies and antibody-like molecules directed against HIV coat proteins, drugs that inhibit viral replication, and cytotoxins targeted
35 to infected cells that express HIV encoded proteins.

- 2 -

Cytotoxic hybrid proteins composed of a cytotoxin fused to part of the CD4 receptor have been proposed as a way to destroy cells expressing HIV encoded proteins. This approach relies on the fact that the HIV envelope protein, gp120, recognizes the CD4 receptor, which is present on T4 lymphocytes and certain cells of the monocyte/macrophage lineage. Thus, a soluble derivative of CD4 might be used to target a cytotoxin to HIV infected cells that express surface gp120. This approach is likely to be most effective against productively infected cells in which HIV is replicating; under these circumstances there is likely to be significant expression of gp120 on the cell's surface. Chaudhary et al. (*Nature* 335:369, 1988) found that administration of a CD4-*Pseudomonas* exotoxin hybrid protein to a lymphocytic cell line chronically infected with HIV causes a decrease in overall protein synthesis. Till et al. (*Science* 242:1166, 1988) found that a CD4-ricin A fusion protein decreases DNA synthesis in cultures of chronically infected H9 cells. In a variation of this strategy, Capon et al. (*Nature* 337:529, 1989) designed a hybrid protein composed of soluble CD4 and the constant region of an antibody. This molecule is designed to direct immune system response to HIV and the HIV coat protein, gp120. Another molecule of this general type has been shown to activate complement (Trauneker et al., *Nature* 339:78, 1989).

Human T-lymphotropic retrovirus type I (HTLV-I) is associated with adult T-cell leukemia and may play a role in other diseases including tropical spastic paraparesis and HTLV-I associated myelopathy.

Epstein-Barr virus (EBV) is a B lymphotropic human herpes virus. The majority of people infected with EBV develop infectious mononucleosis. EBV is also associated with African Burkitt's lymphoma, anaplastic

- 3 -

nasopharyngeal carcinoma, and, in immunocompromised individuals, lymphocytic (usually B-cell) lymphomas.

Summary of the Invention

In general the invention features a method for
5 treating a patient infected with a virus; the method includes administering to the patient a molecule which is capable of specifically binding to a proteinaceous cell receptor expressed on a cell of the patient and which contributes to the disease state of the patient, the
10 molecule being capable of decreasing the viability of the cell. By "cell receptor" is meant a molecule which is encoded by cellular DNA, binds a ligand, and is expressed so that at least a portion of the molecule is exposed on the cell surface. By "specifically binding" is meant
15 that the molecule does not substantially bind to other cell receptors or cell surface proteins. By "reduces viability" is meant kills or interferes with proliferation. By "ligand" is meant a molecule which is capable of binding to a protein.

20 In preferred embodiments, the virus is human immunodeficiency virus; the virus is HTLV-I; the virus is EBV; the proteinaceous cell receptor is the high affinity interleukin-2 receptor; the molecule kills cells bearing the cell receptor; the molecule is a hybrid molecule
25 which includes a first and a second portion joined together covalently, the first portion includes a molecule capable of decreasing cell viability and the second portion includes a molecule capable of specifically binding to the cell receptor. In more
30 preferred embodiments, the second portion of the hybrid molecule includes all or a binding portion of an antibody specific for the cell receptor; the second portion of the hybrid molecule includes all or a binding portion of a ligand for the cell receptor. By a "binding portion" is
35 meant a portion capable of specifically binding to a cell

- 4 -

receptor. In still more preferred embodiments, the ligand is a growth factor; the ligand is an interleukin. In yet more preferred embodiments, the interleukin is interleukin-4; the interleukin is interleukin-6.

- 5 In a preferred embodiment, the first portion of the hybrid molecule includes a cytotoxin. In a more preferred embodiment, the cytotoxin is a fragment of a peptide toxin which is enzymatically active but which does not possess generalized eukaryotic receptor binding
- 10 activity. In an even more preferred embodiment, the fragment of a peptide toxin includes fragment A of diphtheria toxin and enough of fragment B of diphtheria toxin to form a pore in a cell membrane. In still more preferred embodiments, the molecule is DAB₄₈₆IL-2; the
- 15 molecule is DAB₃₈₉IL-4; the molecule is DAB₃₈₉IL-6.

In a preferred embodiment, the molecule includes all or a binding portion of an antibody specific for the cell receptor. In a more preferred embodiment, the antibody is a complement activating antibody.

- 20 The invention provides a method for treating viral diseases. The method eliminates or neutralizes cells which bear a cell surface receptor which is induced following viral infection of the patient; this is accomplished by targeting a molecule (e.g., a cytotoxin
- 25 or lytic antibody) to that receptor. The cells targeted can be either infected or uninfected cells of the patient who is infected with the virus. Viruses can cause infected cells to express a cell surface receptor; in some cases, the viral disease causes some uninfected
- 30 cells to express certain receptors, and this expression may contribute to the pathophysiology of the viral disease. Virally induced expression of a receptor a particular cell includes: expression by a cell (whether infected or uninfected) of a receptor which would
- 35 normally never be expressed by that cell (or cell type);

- 5 -

expression by a cell (whether infected or uninfected) of a receptor which can normally be expressed by that cell (or cell type) but which is normally expressed only under certain circumstances; and virally associated expression of a receptor by a cell which expression is significantly higher than the expression by an unaffected cell. For example, HTLV-I infected T-lymphocytes persistently express the high-affinity interleukin-2 receptor; normally this receptor is expressed by T-lymphocytes only after cell activation. In other instances, including HIV infection itself, productive viral replication does not occur until a circumstantial event triggers or markedly elevates expression of a receptor in association with a state of cellular activation. The method of the invention is applicable to this situation as well. To summarize, virally associated expression of a cell receptor includes: any expression of a cell receptor that is induced, directly or indirectly, by the virus; cell receptor expression which follows infection with the virus; and cell receptor expression which contributes to the disease state.

The method of the invention can be used to treat HIV infection by killing or neutralizing HIV-infected and harmful uninfected cells that express the high affinity interleukin-2 receptor (IL-2R). *In vitro* experiments (see below) have demonstrated that the high affinity IL-2 receptor is expressed on the surface of HIV-infected cells prior to expression of HIV encoded proteins, and the method of the invention is thus useful for destroying HIV infected lymphocytes and monocytes/macrophages both before and after viral replication has occurred. Accordingly, the method of the invention is capable of eliminating HIV-infected cells at an early stage, prior to the production and release of mature virus. The method of the invention can eliminate chronically and

- 6 -

latently infected cells that express the high affinity IL-2R. Because the method of the invention targets a protein encoded by cellular DNA rather than an HIV-encoded protein, the method is useful for treating

5 patients infected with any strain of HIV. HIV is known to undergo rapid mutation particularly in the sequences encoding the coat protein and the reverse transcriptase. The method of the invention will be effective against HIV regardless of the exact structure of the coat protein or

10 the reverse transcriptase and thus should be effective against all HIV variants. This is important because the HIV in a given patient is likely to consist of a cohort of viral species. The method of the invention will be effective against all forms of HIV (e.g., HIV-1 and HIV-

15 2) providing that following infection, expression of a cell surface receptor, such as IL-2R, is induced.

Infection by EBV or HTLV-I is also thought to cause expression of the interleukin-2 receptor on the surfaces of cells which would otherwise not express that

20 receptor. Thus, the method of the invention may be used for treatment of diseases caused by infection with EBV or HTLV-I.

Generally, the method of the invention will be useful for treatment of any viral infection which is

25 associated with the induction of expression of a cell surface receptor. In addition to eliminating or neutralizing virally infected cells, the method of the will be useful for treatment of virally caused diseases in which virally induced expression of a cell surface

30 receptor contributes to the course of the infection or to the pathophysiology of the disease. Although the interleukin-2 receptor is suggested as a target in the examples below, the method of the invention can be used to target cells bearing other virally induced receptors

- 7 -

(e.g., the interleukin-4 receptor or the interleukin-6 receptor).

HIV and Immune Cell Activation

Immune cell activation and accompanying cytokine
5 production play an extremely important role in the progression of HIV infection and in the pathologic effects of HIV infection.

Activation of HIV infected cells has been shown to be important for the establishment of a productive
10 infection. The activation signals which have been shown to be capable of inducing cell activation and virus production include: phorbol esters, UV irradiation, antigens, mitogens (e.g., phytohemagglutinin), and cytokines such as tumor necrosis factor α , tumor necrosis
15 factor β , granulocyte-macrophage colony-stimulating factor, and interleukin-6 (see McCune, *Cell* 64:351, 1991 for a review). Resting T cells, which represent the vast majority of T-cells *in vivo*, can bind and take up HIV; these cells begin to synthesize viral DNA, but they fail
20 to convert HIV genomic RNA to full length double-stranded DNA (Zack et al., *Cell* 61:213, 1990). Thus, infection of resting cells may lead to an eclipsed or latent viral state which can be converted to a productive viral infection only after cell activation. In other cases,
25 the infected cell may be briefly activated, allowing viral integration and render the cell fully permissive for HIV infection. This cell can then cycle back to a resting state, possibly allowing the establishment of a second latent infection. *In vitro* studies have
30 demonstrated that infected cells may remain in a latent state in which the viral genome is integrated, but viral transcription occurs at a very low level (Pomercantz et al., *Cell* 61:1271, 1990). Viral RNA production in such latently infected cells is dramatically increased by cell
35 activation. Cell activation triggers high level

- 8 -

expression of virally encoded proteins and viral replication. High level expression and viral replication leads to a burst of virion release and host cell death. The fact that mitogen- and antigen-induced cell

5 activation induces HIV expression in T-lymphocytes and monocytes/macrophages has led to the hypothesis that cytokines play an important role in the activation of HIV. A protein, NF- κ B, whose expression is induced

10 for the enhanced HIV expression which accompanies cell activation. This protein binds to sequences present in the HIV enhancer and, along with other transcription factors, contributes to the activation of HIV expression (Lenardo et al., *Cell* 58:227, 1989; Böhnlein et al., *Cell*

15 53:827, 1988) and to the activation of IL-2R expression (see below). It appears that normal immune cell activation and accompanying cytokine production may contribute to the progression of HIV infection at several points including: establishment of HIV infection,

20 enhancement of viral protein expression and replication in productively infected cells, and escape from latency. Accordingly, administration of immunostimulants to patients suffering from HIV-induced immunosuppression may, paradoxically, be detrimental since such agents may

25 induce activation and proliferation of resting T-lymphocytes and monocytes/macrophages and thus enhance viral spread.

HIV may itself initiate immune cell activation and interleukin-2 receptor expression. *In vitro* experiments

30 have demonstrated that exposure of CD4+ lymphocytes or CD4+ cells of the monocyte/macrophage lineage to HIV or purified gp120 results in cell activation and expression of the p55 subunit of the high-affinity interleukin-2 receptor, a molecule which is not normally expressed on

35 the surface of resting cells (Kornfeld et al., *Nature*

- 9 -

35:445, 1988; Allen et al. *J. Clin. Invest.* 85:192, 1990). These results are consistent with the observed increase in surface IL-2 receptor on the monocytes of HIV infected patients (Allen et al., *supra*) and with the
5 observed increase in soluble IL-2R in the serum of HIV infected individuals (Prince et al., *J. Immunol.* 140:1139, 1988; Sethi et al., *Immunol. Lett.* 13:179, 1986). This HIV-triggered expression of surface p55 occurs prior to expression of HIV encoded proteins
10 (Kornfeld et al., *supra*; Allen et al. *supra*; Fields et al., *Nature* 333:278, 1988; Wahl et al., *Proc. Natl. Acad. Sci. USA* 86:621, 1989). Since purified gp120 can activate uninfected cells, soluble gp120 may activate latently infected cells and initiate a productive
15 infection. In addition, soluble gp120 may activate uninfected cells thereby producing both a favorable environment for subsequent infection of those cells and more of the cytokines and inflammatory mediators which contribute to the generalized immunological dysfunction
20 observed during the course of AIDS. It is not known to what extent exposure to HIV or soluble gp120 activates lymphocytes and monocytes/macrophages *in vivo*.

The method of the invention can eliminate or neutralize any cell which inappropriately expresses the
25 high affinity IL-2R as a result of HIV infection or contact with HIV or soluble gp120. Resting lymphocytes and monocytes/macrophages do not express the high affinity IL-2R, and thus are unaffected.

Cytokines may also play a more indirect role in
30 the pathology of HIV infection. As described above, exposure of lymphocytes and monocytes/macrophages soluble gp120 may induce immune cell activation (Kornfeld et al., *supra*; Allen et al., *supra*). While these cells are not infected, their activation may adversely affect immune
35 system function. Activated cells undergo premature

- 10 -

differentiation and thus become refractory to activation by a second immunologic stimulus. This phenomenon has been proposed as an explanation for the fact that the many of the immune cells of HIV infected individuals are
5 phenotypically differentiated but functionally impaired (Allen et al., *supra*). Further, the cytokines and inflammatory mediators produced by activated cells may have adverse metabolic and immunological effects.

The importance of cytokines in the progression of
10 HIV infection and in the pathologic effects of HIV infection suggests that it may be possible to treat HIV infection by interfering with cytokine action, for example, by blocking cytokine receptors. Such an approach can interfere with both HIV infection and the
15 pathological effects of excess cytokine expression. This approach can be used on its own or to supplement receptor-targeted cytotoxin mediated elimination or neutralization of cells bearing a virally induced receptor.

20 Described in detail below is an approach to treating HIV infection by targeting molecules to cells bearing the interleukin-2 receptor. However, the method of the invention is not limited to the targeting of cells bearing this receptor. In treatment of HIV infection it
25 may be important to target cells bearing other receptors (e.g., the interleukin-6 receptor or the interleukin-4 receptor). For example, if B-cell hyperactivation contributes to the progression of HIV (Amadori et al., *Immunol Today* 11:374, 1990), drugs which target cytokine
30 receptors present on activated B-cells may indirectly assist in controlling HIV infection. In addition, to the extent that hyperactivation of immune cells contributes to the disruption of normal immune function observed in HIV infection, molecules targeted to activated cells via
35 a cytokine receptor may provide a valuable therapeutic

- 11 -

approach even when the cell targeted is not itself infected. For example, the method of the invention may be useful for treatment of AIDS-related psoriasis. Psoriasis is characterized by the hyperproliferation of epidermal cells, and IL-2R is known to be present in the dermis of psoriatic plaques (Gottlieb et al., *J. Am. Acad. Derm.* 18:6, 1988). Since the epidermis is a part of the immune system and expresses a full complement of cytokines, many of which (e.g., IL-2, TNF and GM-CSF) are thought to play a role in psoriasis (Duvic, *J. Invest. Derm.* 95:38S, 1990), the receptors for these cytokines may provide useful targets according to the method of the invention. The method of the invention will also be useful for treating some neoplasms that arise in the later stages of HIV infection and which involve IL-2R bearing cells (e.g., Kaposi's sarcoma, and AIDS associated lymphomas).

HTLV, EBV and Immune Cell Activation

HTLV-I infection of T-lymphocytes is generally associated with persistent expression of the high affinity interleukin-2 receptor. This is in contrast to the transient expression of the high affinity interleukin-2 observed after normal cell activation. Persistent expression of the high affinity interleukin-2 receptor is thought to be mediated by the HTLV-I tat-I gene product which induces expression of NF- κ B, a powerful transcriptional activator. NF- κ B in turn stimulates the expression of the p55 subunit of the interleukin-2 receptor. This subunit in combination with the p70 subunit, which is constitutively expressed, forms the high affinity interleukin-2 receptor. HTLV-I infected patients have a high degree of spontaneous lymphocyte proliferation, consistent with an antigen-independent increase in interleukin-2 receptor expression. Further, persistent expression of the high

- 12 -

affinity receptor IL-2R may contribute to HTLV-I mediated cell transformation. Infection by HTLV-II, a related retrovirus also causes an increase in antigen-independent lymphocyte proliferation and may also be associated with
5 an increase in interleukin-2 receptor expression. Thus, IL-2R targeted molecules will be useful for treatment of HTLV-I or HTLV-II infection.

As outlined above, NF- κ B plays a role in the pathology of both HIV and HTLV-I expression; For
10 example, the HIV enhancer has two binding sites for NF- κ B, and induction of NF- κ B induces viral transcription. Since NF- κ B is activated by many of the same signals which activate T-cells, including mitogens, phorbol esters, antibodies against cell surface markers which
15 mimic physiologic T cell activation, and activators of protein kinase C, NF- κ B may play an important role in the course of HIV infection. Other viruses, including cytomegalovirus have NF- κ B binding sites within their enhancers. Further, cytomegalovirus and hepatitis virus
20 B, like HTLV-I, encode trans-activators which induce NF- κ B expression. A number of viruses induce NF- κ B expression, and any such virus may also induce expression of interleukin-2 receptor on the surface of cells which they infect. To the extent which expression of the
25 receptor (or some other receptor) is induced, diseases caused by such viruses may be treated by the method of the invention.

EBV infection is known to induce expression of the high-affinity interleukin-2 receptor on infected cells.
30 Accordingly the method of the invention can be used to kill or neutralize EBV infected cells.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

- 13 -

Detailed Description

The drawings will first briefly be described.

Drawings

Fig. 1 is a graphical representation of the effect
5 of DAB₄₈₆IL-2 on viability of both HIV-infected and
uninfected CD4+ T cells. The percentage of viable cells
is presented as a function of the number of days post
infection.

Fig. 2 is a representation of the results of a
10 SDS-PAGE analysis of proteins in HIV-infected and
uninfected CD4+ T cells.

Fig. 3 is a graphical representation of the effect
of DAB₄₈₆IL-2 on reverse transcriptase activity present in
HIV infected and uninfected monocytes. Reverse
15 transcriptase activity (cpm) is presented as a function
of DAB₄₈₆IL-2 concentration (nM) for uninfected (open
bars) and HIV-1 (filled bars) infected cells.

Molecules Useful in the Method of the Invention

In general, there are three ways in which the
20 molecules useful in the invention can act: (1) the
molecule can kill a cell because the molecule has a
cytotoxic domain; (2) the molecule (an antibody) can
cause cell lysis by inducing complement; and (3) the
molecule can block binding or uptake of receptor's
25 ligand. In all three cases the molecule must be targeted
to receptor bearing cells; this is accomplished by
including the receptor's ligand (or a portion or
derivative thereof) or an anti-receptor antibody as part
of the molecule.

30 Interleukin-2 receptor targeted molecules useful
for treatment of HIV infection provide examples of each
of these three approaches. A fusion molecule which
includes the IL-2 receptor binding portion of IL-2 and a
cytotoxin can be used to kill HIV infected activated
35 lymphocytes and monocytes/macrophages. This molecule can

- 14 -

also kill uninfected IL-2 receptor-bearing cells. Such uninfected cells may contribute to the disease state. Similarly, the second type of molecule described above, a complement fixing antibody, in this instance directed
5 against the IL-2 receptor, can eliminate infected and uninfected, IL-2 receptor-bearing cells in a patient infected with HIV. In this example, the third type of molecule could be a molecule that blocks binding of IL-2 to its receptor. This molecule would prevent infected
10 cells from receiving a proliferation signal from IL-2 and thus could slow the spread of HIV infection.

Molecules useful for killing or neutralizing IL-2R bearing cells of HIV infected individuals can take a number of forms. When IL-2 itself is the targeting
15 agent, the molecule can be a cytotoxic hybrid molecule in which IL-2 is fused to a toxin molecule, preferably a polypeptide toxin. Derivatives of IL-2 which bind to IL-2R, lack IL-2 activity and block binding and/or uptake of bona fide IL-2 are useful in the method of the invention
20 because they will prevent IL-2-induced proliferation of IL-2R bearing cells. When an anti-IL-2R antibody is the targeting agent, a cytotoxic hybrid molecule can be formed by fusing all or part of the antibody to a cytotoxin. The effectiveness of such an antibody/toxin
25 hybrid, like that of an IL-2/toxin hybrid, depends on the hybrid molecule being taken up by cells to which it binds. Anti-IL-2R antibodies which block binding and/or uptake of IL-2 are also useful in the method of the invention. Lytic anti-IL-2R antibodies are useful in the
30 invention because they can direct complement against IL-2R-bearing cells and thus cause their lysis.

Some the molecules can be hybrid molecules formed by the fusion of all or part of two or more molecules. The hybrid molecule can be a hybrid protein encoded by a
35 recombinant DNA molecule, in which case the two domains

- 15 -

are joined (directly or through an intermediary domain) by a peptide bond. Alternatively, two domains can be produced separately and joined by a covalent bond in a separate chemical linkage step. In some cases, the
5 cytotoxic domain of a hybrid molecule may itself be derived from two separate molecules.

Interleukin-2 as a Targeting Agent

IL-2 or any IL-2 receptor binding derivative thereof can be used as a targeting agent for a cytotoxin.
10 The DNA and amino acid sequences of IL-2 are known (Tadatsugu et al., *Nature* 302:305, 1983), and its structure has been predicted by x-ray crystallography (Brandhuber et al., *Science* 238:1707, 1987). Analysis of genetically engineered variants of IL-2 has provided some
15 information concerning which residues are important for IL-2R binding (Collins et al., *Proc. Natl. Acad. Sci. USA* 85:7709, 1988) and bioactivity (Cohen et al. *Science* 234:349, 1989; Collins et al., *supra*). Variants of IL-2 which are useful in the invention include deletion
20 mutants (Genbauffe et al., USSN 388,557, hereby incorporated by reference) which lack one or more amino acid residues in the region between residue 74 and residue 79 (numbering according to Williams et al., *Nucl. Acids Res.* 16:1045, 1988). These mutants effectively
25 target toxins to IL-2R-bearing cells (Genbauffe et al., *supra*). Generally, IL-2 variants useful for targeting a cytotoxin must efficiently bind IL-2R and be endocytosed. The ability of various derivatives to bind to the IL-2 receptor can be tested with an IL-2R binding assay
30 described below.

In designing molecules targeted to cells bearing the IL-2 receptor it must be recognized that the IL-2 receptor, like other receptors, has several forms; and it may be desirable to target cells bearing one form and not
35 another. The human interleukin-2 receptor has a high-

- 16 -

an intermediate-, and a low-affinity form. The high affinity receptor has an apparent K_d of $\sim 10^{-10}M$ and is composed of two subunits, p55 and p75 (also called p70). When expressed on the cell surface, both the p75 and p55 subunits are capable of binding IL-2. The p75 subunit corresponds to the intermediate affinity receptor ($K_d \sim 8.2 \times 10^{-10}M$), and p55 subunit corresponds to the low affinity receptor ($K_d \sim 1-3 \times 10^{-8}M$). The p75 subunit is expressed on the surface of resting T cells, natural killer cells monocytes/macrophages, and lymphokine-activated killer (LAK) cell precursors, while the high affinity receptor is expressed on activated T- and B-cells.

In the method of the invention it may be desirable to target only cells bearing the high affinity receptor. In these circumstances useful molecules will eliminate or neutralize cells bearing the high affinity IL-2 receptor at a concentration which leaves cells bearing the intermediate or low affinity receptor largely unaffected. When the molecule, like IL-2 itself, has affinity for all three classes of IL-2 receptor, selectivity can be accomplished by administering the molecule at a concentration which does not permit significant binding to cells bearing lower affinity receptors. A hybrid molecule may have altered receptor affinities compared to IL-2. Such hybrid molecules may be more or less selective for cells bearing the high affinity IL-2 receptor. For example, cells bearing the high-affinity receptor are 500-1000 times more sensitive to DAB₄₈₆IL-2, a fusion protein consisting of part of diphtheria toxin and part of IL-2, than are cells bearing the intermediate- affinity receptor (Waters et al., *Eur. J. Immunol.* 20:785, 1990).

A cytotoxin can be attached to an IL-2 derivative in a number of ways. Preferably, an IL-2/toxin hybrid is

- 17 -

a hybrid protein produced by the expression of a fused gene. Alternatively, the cytotoxin and the IL-2 derivative can be produced separately and later coupled by means of a non-peptide covalent bond. Linkage methods
5 are described below.

Interleukin-4 as a Targeting Agent

Interleukin-4 (IL-4) is a cytokine which acts on a variety of cell types. Its receptor is expressed on a number of cell types, including CD4+ T cells and
10 monocytes. IL-4 can act as a T cell growth factor and it is thought to have an influence on IL-2 induced lymphocyte proliferation.

A cytotoxin directed against IL-4 receptor-bearing cells may enhance the effectiveness of molecules directed
15 against IL-2R-bearing cells. The protein and DNA sequence of IL-4 are known (Lee et al., *J. Biol. Chem.* 263:10817, 1988). IL-4 can be used to create hybrid IL-4/toxin molecules similar to IL-2/toxin hybrid molecules.

20 Monoclonal Antibodies as Targeting Agents

Monoclonal antibodies directed against IL-2R, IL-4R or any cell receptor of choice can be used to direct toxins to cells bearing that receptor. These antibodies or antibody fragments can be fused to a cytotoxin either
25 by virtue of the toxin and the antibody being encoded by a fused gene which encodes a hybrid protein molecule, or by means of a non-peptide covalent bond which is used to join separately produced ligand and toxin molecules. Several useful toxins are described below.

30 Antibody/toxin hybrids can be tested for their ability to kill receptor bearing cells using a toxicity assay similar to that which is described below for IL-2R bearing cells.

Toxins

- 18 -

The toxin molecules useful in the method of the invention are preferably toxins, such as peptide toxins, which are significantly cytotoxic only when present intracellularly. Of course, under these circumstances
5 the molecule must be able to enter a cell bearing the targeted receptor. This ability depends on the nature of the molecule and the nature of the cell receptor. For example, cell receptors which naturally allow uptake of a ligand are likely to provide a means for a molecule which
10 includes a toxin to enter a cell bearing that receptor. Preferably, a peptide toxin is fused to an IL-2R binding domain by producing a recombinant DNA molecule which encodes a hybrid protein molecule. Such an approach ensures consistency of composition.

15 Many peptide toxins have a generalized eukaryotic receptor binding domain; in these instances the toxin must be modified to prevent intoxication of non-receptor bearing cells. Any such modifications must be made in a manner which preserves the cytotoxic functions of the
20 molecule (see U.S. Department of Health and Human Services, U.S. Serial No. 401,412). Potentially useful toxins include, but are not limited to: cholera toxin, ricin, 0-Shiga-like toxin (SLT-I, SLT-II, SLT II_v), LT toxin, C3 toxin, Shiga toxin, pertussis toxin, tetanus
25 toxin, *Pseudomonas* exotoxin, alorin, saporin, modeccin, and gelanin.

Diphtheria Toxin-based Molecules

Diphtheria toxin can be used to produce molecules useful in the method of the invention. Diphtheria toxin,
30 whose sequence is known, is described in detail in Murphy U.S. Patent 4,675,382, hereby incorporated by reference. The natural diphtheria toxin molecule secreted by *Corynebacterium diphtheriae* consists of several functional domains which can be characterized, starting
35 at the amino terminal end of the molecule, as

- 19 -

enzymatically-active Fragment A (amino acids Gly₁ - Arg₁₉₃) and Fragment B (amino acids Ser₁₉₄ - Ser₅₃₅), which includes a translocation domain and a generalized cell binding domain (amino acid residues 475 through 535).

5 The process by which diphtheria toxin intoxicates sensitive eukaryotic cells involves at least the following steps: (i) the binding domain of diphtheria toxin binds to specific receptors on the surface of a sensitive cell; (ii) while bound to its receptor, the
10 toxin molecule is internalized into an endocytic vesicle; (iii) either prior to internalization, or within the endocytic vesicle, the toxin molecule undergoes a proteolytic cleavage between fragments A and B; (iv) as the pH of the endocytic vesicle decreases to below 6, the
15 toxin crosses the endosomal membrane, facilitating the delivery of Fragment A into the cytosol; (v) the catalytic activity of Fragment A (i.e., the nicotinamide adenine dinucleotide - dependent adenosine diphosphate (ADP) ribosylation of the eukaryotic protein synthesis
20 factor termed "Elongation Factor 2") causes the death of the intoxicated cell. It is apparent that a single molecule of Fragment A introduced into the cytosol is sufficient to block down the cell's protein synthesis machinery and kill the cell. The mechanism of cell
25 killing by *Pseudomonas* exotoxin A, and possibly by certain other naturally-occurring toxins, is very similar.

DAB₄₈₆IL-2, a fusion protein in which the receptor binding domain of diphtheria toxin has been replaced by a
30 portion of human IL-2 (Williams et al., *J. Biol. Chem.* 35:20673, 1990; see also Williams et al., *Protein Eng.* 1:493, 1987), is an example of a molecule useful in the method of the invention. This molecule selectively kills IL-2R-expressing tumor cells and lymphocytes (Waters et
35 al., *Eur. J. Immunol.* 20:785, 1990; Kiyokawa et al.,

- 20 -

Cancer Res. 49:4042, 1989). Because of its ability to kill activated lymphocytes, DAB₄₈₆IL-2 has been used to control graft rejection (Pankewycz et al., *Transplantation* 47:318, 1989; Kickman et al., *Transplantation* 47:327, 1989) and to treat certain autoimmune disorders (Forte et al., *2nd International Symposium on Immunotoxins*, 1990).

DAB₄₈₆IL-2 is a chimeric molecule consisting of Met followed by amino acid residues 1 through 485 of the mature diphtheria toxin fused to amino acid residues 2 through 133 of IL-2. Thus, DAB₄₈₆IL-2 includes all of diphtheria toxin fragment A, which encodes the enzymatically active portion of the molecule, and a portion of fragment B. The portion of fragment B present in DAB₄₈₆IL-2 does not include the generalized receptor binding domain but does include the translocation domain which facilitates delivery of the enzymatically active portion into the cytosol.

Experimental Methods

Presented below are experiments which demonstrate that DAB₄₈₆IL-2 can kill HIV-1 infected T cells, selectively eliminate HIV-1 infected cells from mixed cultures of infected and uninfected T cells, and reduce HIV-1 replication in cultures of infected monocytes. DAB₄₈₆IL-2 is also shown to inhibit production of viral proteins in cultures of infected T cells and to block production of infectious HIV-1 in cultures of infected T cells.

Other molecules targeted to IL-2R may be screened using the methods described below.

Preparation of DAB₄₈₆IL-2

DAB₄₈₆IL-2 was produced in *E. coli* harboring the DAB₄₈₆IL-2 encoding plasmid, pDW24 (Williams et al., J. Biol. Chem. 265:20673, 1990, except amp^r is replaced by kan^r). The protein was purified by immunoaffinity

- 21 -

chromatography and high pressure liquid chromatography (Williams et al., supra).

Killing of HIV-1 Infected T Cells by DAB₄₈₆IL-2

DAB₄₈₆IL-2 at 10^{-8} M or 10^{-9} M destroyed HIV-1
5 infected T cells while having almost no effect on uninfected T cells. CD4+ T cells were prepared from the peripheral blood of HIV-1 negative donors by negative selection to remove B cells, macrophages, natural killer cells, and CD8+ T cells. B cells and macrophages were
10 removed by passage over glass wool. CD8+ and CD16+ cells were removed using anti-CD8 (OKT8, American Type Culture Collection, Rockville, MD) and anti-CD16 (Leu 11a, Bectin-Dickinson, Mountain View, CA) monoclonal antibody coated magnetic beads by the method of Haregewoin et al.
15 (Nature 340:309, 1989). Cells were cultured at 2×10^6 /ml in RPMI 1640 (GIBCO/BRL, Bethesda, MD) and 10% bovine calf serum (GIBCO/BRL, Bethesda, MD) supplemented with lymphocult-T (Boehringer-Mannheim Biochemicals, Indianapolis, IN; corresponds to $\sim 10^{-9}$ M IL-2). Infections
20 were performed by incubation with HTLVIII_B (prepared from filtered supernatants of infected H9 cells, AIDS Research and Reference Reagent Program NIAID, National Institutes of Health, Bethesda, MD) for 1 hr at a multiplicity of infection of 10 (calibrated by limiting dilution of H9
25 cells). DAB₄₈₆IL-2 (10^{-7} M or 10^{-8} M) was added on days one and three post-infection. Cell cultures were split twice weekly, and viability was determined by a trypan blue dye exclusion assay (Kruse et al., eds. *Tissue Culture: Methods and Applications*, Academic Press, 1989)
30 Referring to FIG. 1, treatment of uninfected cells with 10^{-8} M (open circles) or 10^{-7} M (open squares) DAB₄₈₆IL-2 only transiently impaired proliferation of T cells in response to 10^{-9} M IL-2. Further, cultures of uninfected T cells treated with DAB₄₈₆IL-2 and untreated
35 uninfected T cells achieve similar cell densities after

- 22 -

two weeks of culturing. In contrast, HIV-1 infected cells were eliminated by incubation with 10^{-8} M (filled circles) or 10^{-7} M (filled squares) DAB₄₈₆IL-2. This toxic effect occurs despite the fact that the cells are cultured in the presence of 10^{-9} M IL-2 which is required for cell viability and which has a 10- to 100- fold higher affinity for IL-2R than does DAB₄₈₆IL-2 (Waters et al., *supra*). Infected cells which were not treated with DAB₄₈₆IL-2 were >75% viable after two weeks, demonstrating that the reduction in viability of DAB₄₈₆IL-2 exposed cells is a specific effect and is not due to viral replication.

Selective Killing of HIV-1 Infected T Cell by DAB₄₈₆IL-2

DAB₄₈₆IL-2 prevented production of the HIV-1 encoded proteins, gp120, p55, and p24, in mixed cultures of HIV-1 infected and uninfected T cells.

CD4+ T cells prepared and infected as described above were incubated overnight in RPMI 1640 medium containing 10% bovine calf serum and washed three times with the same medium prior to addition of uninfected cells from the same donor. Infected T cells (10^9) and uninfected T cells (10^7) were cultured in the presence of 10^{-9} M IL-2 (Lymphocult T) with or without 10^{-8} M DAB₄₈₆IL-2 at 10^6 cells/ml. Every two days cells were pelleted, washed twice, and resuspended in fresh IL-2 containing media. DAB₄₈₆IL-2 was added to the treated cultures on days 1 and 3 and was washed out 24 hours later. The untreated cultures were washed according to the same schedule but were not exposed to DAB₄₈₆IL-2. Two weeks after infection, the cells were labelled overnight with ³⁵S in methionine-free media (GIBCO/BRL, Bethesda, MD). Proteins were immunoprecipitated (Coligan et al, eds., *Current Protocols in Immunology*, John Wiley & Sons, New York, 1991) with either anti-HIV globulin (NIAID) or the

- 23 -

framework anti-MHC class I antibody (W6/32).

Immunoprecipitated proteins were separated on SDS gels and visualized by autoradiography.

Referring to FIG. 2, the proteins in lanes 2-4 were immunoprecipitated using anti-HIV antibody; the proteins in lanes 6-9 were immunoprecipitated using anti-MHC class I antibody; lane 1 has size markers; and the arrows along the right side indicate the expected positions of HIV-1 encoded proteins gp120, p55 and p24. The HIV-1 encoded proteins gp160, p55 and p24 were present in untreated mixed culture (lane 4), but could not be detected in the DAB₄₈₆IL-2 treated culture (lane 5). As expected, normal MHC class I proteins were detected in both the treated (lane 9) and untreated (lane 8) mixed T cell cultures. Unmixed cultures of both uninfected cells (lanes 2 and 6) and infected cells (lanes 3 and 7) served as controls. Cell viability was >95% for both treated and untreated mixed cultures, indicating that the cytotoxic action of DAB₄₈₆IL-2 was limited to infected cells.

DAB₄₈₆IL-2 treatment of mixed cultures of infected and uninfected T cells completely eliminated production of HIV-1 p24 protein. This result was demonstrated by means of a sensitive ELISA assay.

Treated and untreated mixed cultures of infected and uninfected CD4+ T cells were prepared as described above. Cells were pelleted and resuspended twice weekly, and the culture supernatants were assayed for the presence of p24 two days later using an ELISA assay (Abbott, Chicago, IL).

Referring to Table 1, in untreated mixed cultures, the level of p24 increased steadily for at least 18 days post-infection. In contrast, p24 was undetectable in mixed cultures which were treated with 10^{-8} M DAB₄₈₆IL-2.

- 24 -

This experiment demonstrates that DAB₄₈₆IL-2 can block all production of an HIV-1 encoded protein.

Table 1:

Incubation of Mixed T Cell Cultures with DAB₄₈₆IL-2

5 Eliminates Production of p24 Antigen

	Days Cultured	Infected T-Cells	HIV-1 Infected T-Cells	Mixed T Cell Cultures	Mixed T Cell Cultures + DAB ₄₈₆ IL-2 ²
10	6	0	0	0	0
	9	0	256 ¹	0	0
	12	0	>500	89	0
	15	0	>500	298	0
	18	0	>500	>500	0

15

¹ Picograms of p24 in the supernatant two days after cell washing.

² 10⁻⁸M DAB₄₈₆IL-2 add on days 1 and 3.

DAB₄₈₆IL-2 Prevents HIV-1 Infection

Treatment of mixed cultures of infected and uninfected T cell with DAB₄₈₆IL-2 prevented the production of infectious HIV. This was demonstrated by comparing the p24 produced in a co-culture of H9 tumor cells and cells from a DAB₄₈₆IL-2 treated mixed infected and uninfected T cell culture with that produced by a co-culture of H9 tumor cells and cells from an untreated mixed infected and uninfected T cell culture. Despite the fact that H9 cells can be readily infected by HIV, no HIV p24 could be detected in the co-culture supernatant if the mixed infected and uninfected T cells had been treated with DAB₄₈₆IL-2 at least three days prior to their addition to H9 cells.

CD4+ T cells were purified, infected with HIV-1, washed, and then mixed with uninfected cells at a ratio of 1:10 as described above. Cells were cultured in 10⁻⁹M

- 25 -

IL-2 with or without 10^{-8} M DAB₄₈₆IL-2 (added on days 1 and 3). On days 0, 1, 6 and 9, 5×10^5 T cells were collected, washed and co-cultured with 5×10^5 uninfected H9 tumor cells (NIAID). Co-cultures were incubated for 6 days prior to measurement of the p24 present in the co-culture supernatant. p24 was measured by the ELISA assay described above.

Table 2:Incubation of T Cell Cultures with DAB₄₈₆IL-2

10 Eliminates Infectious Virus

15	Cell Culture	DAB ₄₈₆ IL-2	Day of T cell culture post HIV-1 infection	p24 ¹ day 6 of co-culture
	Uninfected T Cells	----	0	
20	HIV-1 infected T cells	----	0	>500
	T Cell mixture	----	0	>500
	T Cell mixture	10^{-8} M	0	>500
	T Cell mixture	----	1	>500
	T Cell mixture	10^{-8} M	1	>500
25	T Cell mixture	----	6	>500
	T Cell mixture	10^{-8} M	6	0
	T Cell mixture	----	9	>500
	T Cell mixture	10^{-8} M	9	0

30

¹ Measured in picograms

Referring to Table 2, when the mixed infected and uninfected T cell culture was not treated with DAB₄₈₆IL-2, p24 was produced in the co-culture supernatant. This result implies that infectious HIV was produced in the untreated mixed cell culture. In contrast, when cells from a DAB₄₈₆IL-2 treated mixed infected and uninfected T cell culture were added to H9 tumor cells 3 or 9 days

- 26 -

after the second addition of DAB₄₈₆IL-2, no p24 could be detected in the co-culture supernatant. This result implies that DAB₄₈₆IL-2 treatment was able to clear the mixed infected and uninfected cell culture of infectious HIV. DAB₄₈₆IL-2 treated cultures responded normally to IL-2 and phytohemagglutinin indicating that the treatment is not generally toxic to T cells.

DAB₄₈₆IL-2 Eliminates HIV-1 Replication in Monocytes

Treatment with DAB₄₈₆IL-2 blocked HIV-1 replication in monocytes as judged by a reverse transcriptase assay. Aliquots of monocytes (10^7) purified according to Wahl et al. (*Cell Immunol.* 85:3553, 1984) were suspended in 1 ml of primary macrophage culture supernatant containing HIV-1/HTLV-III_{Ba-L} (2.5×10^4 cpm reverse transcriptase activity). As a control, aliquots of monocytes were suspended in RPMI 1640 containing 10% fetal calf serum (FCS, GIBCO/BRL). Infected and control cultures were incubated for 1 h at 37°C with intermittent gentle agitation. The cells were then washed with RPMI 1640 plus 10% FCS, resuspended in RPMI 1640 containing 10% FCS, antibiotics, and glutamine, plated (2×10^6) on chamber slides and incubated at 37°C. DAB₄₈₆IL-2 (10^{-8} to 10^{-10} M) was added beginning on day 3 (when IL-2R is expected to be present on infected monocytes) and every third day thereafter. Supernatants from the infected and control monocyte cultures were assayed directly for reverse transcriptase activity by a modification of the method of Spira et al. (*J. Clin. Microbiol.* 25:97, 1987). Briefly, 15 μ l aliquots of supernatant were removed from each culture, 5 μ l of a buffer containing 30% glycerol and 0.5% Triton x-100 was added to each aliquot in order to solubilize any virus present. 25 μ l of a reaction mix containing 5 μ Ci of ³H-thymidine triphosphate (20 Ci/mmol; Du Pont NEN, Boston, MA), 0.45 μ g poly rA oligo (12-18 bases long; Pharmacia,

- 27 -

Piscataway, NJ), and 10mM MgCl₂ was added to each test aliquot. The reactions were incubated at 37°C for 2.5 h, and the reaction products were ethanol precipitated.

Referring to FIG. 3, 10⁻⁸M DAB₄₈₆IL-2 virtually
5 eliminated reverse transcriptase activity in HIV-1
infected monocytes (solid bars). Lower DAB₄₈₆IL-2
concentrations inhibited reverse transcriptase activity
to a lesser extent. DAB₄₈₆IL-2 did not effect the
background measurement of reverse transcriptase activity
10 in uninfected cells (open bars). Further, adherent
monocyte/macrophages remaining in the DAB₄₈₆IL-2 treated
cultures were viable and morphologically
indistinguishable from untreated cells.

Use for DAB₄₈₆IL-2 for Treatment of Kaposi's Sarcoma

15 Certain epidermoid cancers and sarcomas can also
express functional IL-2 receptors. Accordingly,
molecules targeted to cells bearing the IL-2 receptor can
be used for treatment of cancers and sarcomas associated
with viral diseases. IL-2 receptors were demonstrated on
20 Kaposi's sarcoma cells using B-D phycoerythrin labelled
monoclonal anti-TAC (anti-p55) antibody. Two patients
with advanced AIDS and disseminated, chemotherapy
resistant Kaposi's sarcoma underwent one course of 5
doses of DAB₄₈₆IL-2 administered as a daily 90 min
25 intravenous infusion. Both patients had an approximately
30% regression in their skin lesions.

Preparation of DAB₃₈₉IL-4

A synthetic gene encoding human interleukin-4 was
synthesized (Milligen/Bioscience 7500 DNA synthesizer).
30 The IL-4 sequence (Yodota et al., *Proc Nat'l Acad Sci.*
USA, 83:58994, 1986) was modified to incorporate E. coli-
preferred codon usage (deBoer et al., in *Maximizing Gene*
Expression, Reznikioff et al., eds., 1986, Butterworths,
Boston), and restriction endonuclease cleavage sites were
35 added to facilitate subsequent cloning steps. IL-4

- 28 -

coding sequence (His¹ to Ser¹²⁹) was inserted into pDW27 plasmid. pDW27 is derived from pDW24 (Williams et al., J. Biol. Chem. 265:11885, 1990) by deleting DNA corresponding to amino acids 388 to 485 of native diphtheria toxin.

Cytotoxicity of DAB₃₈₉IL-4

The ability of DAB₃₈₉IL-4 to reduce viability of various cell types was measured using an inhibition of protein synthesis assay; the results of this assay are presented in Table 3. IC₅₀ (M) is the concentration of DAB₃₈₉IL-4 required for a 50% decrease in protein synthesis. The rat, Con A-activated, normal splenic lymphocytes were far less sensitive to DAB₃₈₉IL-4 than any of the other cells or cell lines. Since the rat interleukin-4 receptor does not bind human interleukin-4, this result demonstrates the specificity of DAB₃₈₉IL-4. These rat cells are sensitive to a diphtheria toxin/rat interleukin-2 hybrid molecule.

Table 3:

DAB₃₈₉IL-4 Sensitivity of Normal and Neoplastic Cells and Cell Lines

Cell or Cell Line	Classification	IC ₅₀ (M)
T cell origin HUT 102/6TG C91/PL	Human, CTCL, HTLV-I ⁺ Human, HTLV-I ⁺ , transformed	2.9 x 10 ⁻¹¹ 6.3 x 10 ⁻¹¹
B cell origin Raji	Human, Burkitt's lymphoma EBV ⁺	7.2 x 10 ⁻¹⁰
Myelomononuclear cell U937	Human, histiocytic lymphoma	2.0 x 10 ⁻⁹
Normal PBMC PHA activated T cells	Human	1.6 x 10 ⁻¹⁰
Non-primate Con A-activated normal splenic T cells	Rat	>10 ⁻⁷

- 29 -

Preparation of DAB₃₈₉IL-6

A synthetic gene encoding human interleukin-6 was synthesized (Milligen/Bioscience 7500 DNA synthesizer). The IL-6 sequence (Revel et al., EPA 8611404.9) was
5 modified to incorporate *E. Coli* preferred codon usage (deBoer et al., *supra*), and restriction endonuclease cleavage sites were added to facilitate subsequent cloning steps. The entire IL-6 coding sequence was inserted into pDW27 plasmid as described above for
10 DAB₃₈₉IL-4.

Mixed Toxins

The cytotoxic portion of some molecules useful in the invention can be provided by a mixed toxin molecule. A mixed toxin molecule is a molecule derived from two
15 different polypeptide toxins. Generally, as discussed above in connection with diphtheria toxin, polypeptide toxins have, in addition to the domain responsible for generalized eukaryotic cell binding, an enzymatically active domain and a translocation domain. The binding
20 and translocation domains are required for cell recognition and toxin entry respectively. The enzymatically active domain is the domain responsible for cytotoxic activity once the molecule is inside a cell.

Naturally-occurring proteins which are known to
25 have a translocation domain include diphtheria toxin, *Pseudomonas* exotoxin A, and possibly other peptide toxins. The translocation domains of diphtheria toxin and *Pseudomonas* exotoxin A are well characterized (see, e.g., Hoch et al., *Proc. Natl. Acad. Sci. USA* 82:1692,
30 1985; Colombatti et al., *J. Biol. Chem.* 261:3030, 1986; and Deleers et al., *FEBS Lett.* 160:82, 1983), and the existence and location of such a domain in other molecules may be determined by methods such as those

- 30 -

employed by Hwang et al. *Cell* 48:129, 1987); and Gray et al. *Proc. Natl. Acad. Sci. USA* 81:2645, 1984).

One useful IL-2/mixed toxin hybrid molecule is formed by fusing the enzymatically active A subunit of *E. coli* Shiga-like toxin (Calderwood et al., *Proc. Natl. Acad. Sci. USA* 84:4364, 1987) to the translocation domain (amino acid residues 202 through 460) of diphtheria toxin, and to IL-2. This three-part hybrid molecule, SLT-A/DTB'/IL-2, is useful in the method of the invention in the same way as DAB₄₈₆IL-2 described above. The IL-2 portion of the three-part hybrid causes the molecule to attach specifically to IL-2R-bearing cells, and the diphtheria toxin translocation portion acts to insert the enzymatically active A subunit of the Shiga-like toxin into the targeted cell. The enzymatically active portion of Shiga-like toxin, like diphtheria toxin, acts on the protein synthesis machinery of the cell to prevent protein synthesis, thus killing the cell. The difference between these two types of hybrid toxins is the nature of their enzymatic activities: the enzymatic portion of DAB₄₈₆IL-2 catalyzes the ADP-ribosylation by nicotinamide adenine dinucleotide of Elongation Factor 2, thereby inactivating this factor which is necessary for protein synthesis, while the enzymatic portion of SLT-A/DTB'/IL-2 is a ribonuclease capable of cleaving ribosomal RNA at a critical site, thereby inactivating the ribosome. SLT-A/DTB'/IL-2 hybrid would therefore be useful as a treatment for the same indications as DAB₄₈₆IL-2, and could be substituted or used in conjunction with it if, for example, a patient's activated T-cells develop a resistance to DAB₄₈₆IL-2.

Linkage of Toxins to Binding Ligands

The binding ligand and the cytotoxin of useful hybrid molecules can be linked in several ways. If the

- 31 -

hybrid molecule is produced by expression of a fused gene, a peptide bond serves as the link between the cytotoxin and the binding ligand. Alternatively, the toxin and the binding ligand can be produced separately
5 and later coupled by means of a non-peptide covalent bond.

For example, the covalent linkage may take the form of a disulfide bond. In this case, if the binding ligand is a protein, e.g., IL-2, the DNA encoding IL-2
10 can be engineered to contain an extra cysteine codon as described in Murphy et al. U.S. Serial No. 313,599, hereby incorporated by reference. The cysteine must be positioned so as to not interfere with the IL-2R binding activity of the molecule. For example, the cysteine
15 codon can be inserted just upstream of the DNA encoding Pro₂ of the mature form of IL-2. The toxin molecule must be derivatized with a sulfhydryl group reactive with the cysteine the modified IL-2. In the case of a peptide toxin this can be accomplished by inserting a cysteine
20 codon into the DNA sequence encoding the toxin. Alternatively, a sulfhydryl group, either by itself or as part of a cysteine residue, can be introduced using solid phase polypeptide techniques. For example, the introduction of sulfhydryl groups into peptides is
25 described by Hiskey (*Peptides* 3:137, 1981). Derivatization can also be carried out according to the method described for the derivatization of a peptide hormone in Bacha et al. U.S. Patent No. 4,468,382, hereby incorporated by reference. The introduction of
30 sulfhydryl groups into proteins is described in Maasen et al. (*Eur. J. Biochem.* 134:32, 1983). Once the correct sulfhydryl groups are present, the cytotoxin and IL-2R binding ligand are purified, both sulfur groups are reduced; cytotoxin and ligand are mixed;, (in a ratio of
35 about 1:5 to 1:20) and disulfide bond formation is

- 32 -

allowed to proceed to completion (generally 20 to 30 minutes) at room temperature. The mixture is then dialyzed against phosphate buffered saline to remove unreacted ligand and toxin molecules. Sephadex chromatography or the like is then carried out to separate on the basis of size the desired toxin-ligand conjugates from toxin-toxin and ligand-ligand conjugates.

Assays for IL-2 Receptor Binding and IL-4 Receptor Binding

10 The IL-2R binding ability of various molecules can be measured using an IL-2R assay for high affinity (Ju et al., *J. Biol. Chem.* 262:5723, 1987) or intermediate affinity receptors (Rob et al., *Proc. Natl. Acad. Sci. USA* 84:2002, 1987). The IL-4R binding activity of
15 various molecules can be measured using the assay described by Park et al. (*J. Exp. Med.* 166:176, 1984) or the assay described by Foxwell et al. (*Eur. J. Immunol.* 19:1637, 1989).

Assays for Toxicity

20 Molecules of the invention (both antibodies and hybrid molecules) can be screened for the ability to decrease viability of cells bearing the targeted receptor by means of assays such as those described below.

 Toxicity towards IL-2R bearing cells can be tested
25 as follows. Cultured HUT 102/6TG (Tsuda et al., *Proc. Natl. Acad. Sci. USA* 83:9694, 1986) or YT2C2 (Teshigawari et al., *J. Exp. Med.* 165:223, 1987) cells are maintained in RPMI 1640 medium (Gibco, Grand Island, NY) supplemented with 25 mM HEPES (pH 7.4), 2mM l-glutamine,
30 100 U/ml penicillin, 100 µg/ml streptomycin, and 10% fetal calf serum (Hazelton, Lenexa, KS). Cells are seeded in 96-well V-bottomed plates (Linbro-Flow Laboratories, McLean, VA) at a concentration of 1×10^5 per well in complete medium. Putative toxins are added
35 to varying concentrations ($10^{-12}M$ to $10^{-6}M$) and the

- 33 -

cultures are incubated for 18 hrs. at 37°C in a 5% CO₂ atmosphere. Following incubation, the plates are centrifuged for 5 min. at 170 x g, and the medium removed and replaced with 100 µl leucine-free medium (MEM, Gibco) containing 8 µCi/ml (³H-leucine; New England Nuclear, Boston, MA). After an additional 90 min. at 37°C, the plates are centrifuged for 5 min. at 170 x g, the medium is removed, and the cells are collected on glass fiber filters using a cell harvester (Skatron, Sterling, VA). Filters are washed, dried, and counted according to standard methods. Cells cultured with medium alone serve as the control.

Toxicity towards cells bearing IL-4R may be tested by an assay similar to that described above for IL-2R bearing cells, but utilizing MLA144 cells (Rabin et al. *J. Immunol.* 127:1852, 1981) or HUT 102/6TG cells, seeded at 1 x 10⁵ cells per well and incubated for 40 hours.

Therapy

Generally, the molecules of the invention will be administered by intravenous infusion. They may also be administered subcutaneously. Dosages of molecules useful in the methods of the invention will vary, depending on factors such as whether the substance is a cytotoxin, a lytic antibody, or an cell receptor blocking molecule. In the case of toxic molecules the extent of cell uptake is an important factor; less permeable molecules must be administered at a higher dose.

More than 60 patients have received DAB₄₈₆IL-2 in Phase I/II clinical protocols. The molecule is well tolerated with the maximum tolerated dose (MTD) established by transient asymptomatic hepatic transaminase elevations in about 30% of patients treated at the MTD. Anti-tumor effects have been seen in approximately 40% of patients; responses were noted in B-cell leukemias and lymphomas, cutaneous T-cell lymphoma

- 34 -

and Hodgkin's disease (LeMaistre et al., *Blood* 360a:abstract 1429, 1990; Woodworth et al., *Fourth International Conference on Human Retrovirology*, 1991). Serum concentrations of 10^{-8} M DAB₄₈₆IL-2 have been safely
5 achieved in patients with IL-2 receptor expressing malignancies. Significant anti-tumor effects have been observed in highly refractory leukemia/lymphoma patients and these effects have occurred despite the presence of elevated soluble IL-2R levels in all patients. This
10 observation is consistent with data which suggest that soluble IL-2R does not interfere with binding of IL-2 to the high affinity interleukin-2 receptor. Animal and human studies have demonstrated that DAB₄₈₆IL-2 has no general immunosuppressive effect (LeMaistre et al.,
15 *supra*; Woodworth et al., *supra*).

Experiments indicate that binding and internalization of DAB₄₈₆IL-2 by cells bearing the high affinity IL-2 receptor occurs within 30 minutes of exposure, resulting in maximal inhibition of protein
20 synthesis within several hours. Therefore, the molecule should be effective even if the serum half-life is rather short.

For DAB₄₈₆IL-2 a typical course of therapy might be 0.025 to 0.3 mg/kg/day for 10 to 30 days. This course
25 of treatment can be repeated several times to provide effective therapy.

Other Embodiments

The molecules described above act to decrease cell viability by directing a cytotoxin (or a lytic antibody)
30 to a targeted cell. Also useful in the method of the invention are molecules which interfere with the targeted cell's ability to utilize a cytokine.

Derivatives of IL-2 or other cytokines which block utilization of endogenous cytokine are useful for

- 35 -

preventing proliferation of targeted cells. For example, activated cells deprived of IL-2 fail to proliferate and, in the absence of the essential anabolic stimulus provided by IL-2, will eventually die. With regard to

5 HIV infection, if utilization of IL-2 is prevented, the infected process will be disrupted. The ability of a given IL-2 derivative to interfere with IL-2 function can be tested in an IL-2 bioactivity assay such as the one described by Ju et al. (*J. Biol. Chem.* 262:5723, 1987).

10 Hybrid molecules in which the toxin has been rendered inactive can be also used to block a cytokine receptor. A non-toxic mutant diphtheria toxin molecule has been described (Uchida et al. *J. Biol. Chem.* 248:3838, 1973), and this molecule can be used to produce a non-toxic IL-

15 2/diphtheria toxin hybrid. See Svrluga et al. U.S. Serial No. 590,113, hereby incorporated by reference, for an example of such a hybrid molecule.

Monoclonal antibodies can be used to kill or neutralize cytokine receptor-bearing cells in a number of

20 ways. As described above, anti-cytokine receptor antibodies fused to a toxin molecule can be used to deliver the toxin to receptor-bearing cells. Lytic anti-cytokine receptor antibodies can themselves kill cytokine receptor-bearing cells by activating complement. For

25 example, monoclonal antibodies which activate complement can be used to destroy IL-2R-bearing cells. Complement inducing antibodies are generally those of the IgG1, IgG2, IgG3, and IgM isotypes. Monoclonal anti-IL-2R antibodies can be screened for those able to activate

30 complement using a complement-dependent cytotoxicity test, as follows.

Human T-lymphocytes and EBV transformed B-lymphocytes are labeled with ⁵¹Cr sodium chromate and used as target cells; these cells are incubated with hybridoma

35 culture supernatants and with complement, and then the

- 36 -

supernatants are collected and counted with a gamma counter. Those supernatants exhibiting toxicity against activated T-lymphocytes, but not resting T- or B-lymphocytes, are selected (described in detail in by
5 Leonard et al., *Proc. Natl. Acad. Sci. USA* 80:6957, 1983). The desired anti-IL-2 receptor antibody is purified from the supernatants using conventional methods. The specificity of the antibody can be demonstrated by showing that the activity is blocked by
10 exogenous IL-2.

Also useful are antibodies which block binding and/or uptake of a cytokine. For example, monoclonal antibodies which interfere with the binding and/or uptake of IL-2 are useful in the method of the invention because
15 IL-2R bearing cells deprived of IL-2 fail to proliferate. Blocking monoclonal antibodies (and other blocking molecules) can be tested for their ability to interfere with IL-2 bioactivity using the method of Ju et al., (*supra*). Generally, assays useful for blocking
20 molecules will be competitive binding assays which measure the ability of the molecule being to interfere with binding of one or more of the receptor's natural ligands.

Monoclonal antibodies useful in the method of the
25 invention can be made by immunizing mice with human IL-2R⁺ T-lymphocytes, fusing the murine splenocytes with appropriate myeloma cells, and screening the antibodies produced by the resultant hybridoma lines for the requisite IL-2R binding properties by means of an ELISA
30 assay. Antibody production and screening can be performed according to Uchiyama et al. (*J. Immunol.* 126:1393, 1981). Alternatively, useful antibodies may be isolated from a combinatorial library produced by the method of Huse et al. (*Science* 246:1275, 1989).

- 37 -

The invention can employ not only intact monoclonal or polyclonal antibodies, but also an immunologically-active antibody fragment, for example, a Fab or (Fab)₂ fragment; an antibody heavy chain, an
5 antibody light chain; a genetically engineered single-chain Fv molecule (Ladner et al., U.S. Patent No. 4,946,778); or a chimeric antibody, for example, an antibody which contains the binding specificity of a murine antibody, but in which the remaining portions are
10 of human origin.

What is claimed is:

- 38 -

Claims

1 1. A method for preparing a medicament for the
2 treatment of a viral infection, said method comprising
3 combining, with a pharmaceutically acceptable carrier
4 substance, a molecule which is capable of specifically
5 binding to a proteinaceous cell receptor expressed on a
6 cell which contributes to the pathology of said viral
7 infection, said molecule being capable of decreasing the
8 viability of said cell.

1 2. The method of claim 1 wherein said virus is
2 human immunodeficiency virus.

1 3. The method of claim 1 wherein said virus is
2 HTLV-I.

1 4. The method of claim 1 wherein said virus is
2 EBV.

1 5. The method of claim 1 wherein said
2 proteinaceous cell receptor is the high affinity
3 interleukin-2 receptor.

1 6. The method of claim 1 wherein said molecule is
2 a hybrid molecule comprising a first and a second portion
3 joined together covalently, said first portion comprising
4 a molecule capable of decreasing cell viability and said
5 second portion comprising a molecule capable of
6 specifically binding to said cell receptor.

1 7. The method of claim 6 wherein said second
2 portion comprises all or a binding portion of a ligand
3 for said cell receptor.

- 39 -

1 8. The method of claim 7 wherein said ligand is
2 an interleukin.

1 9. The method of claim 6 wherein said first
2 portion comprises a cytotoxin.

1 10. The method of claim 9 wherein said cytotoxin
2 is a fragment of a peptide toxin which is enzymatically
3 active but which does not possess generalized eukaryotic
4 receptor binding activity.

1 11. The method of claim 10 wherein said fragment
2 of a peptide toxin comprises fragment A of diphtheria
3 toxin and enough of fragment B of diphtheria toxin to
4 form a pore in a cell membrane.

1 12. The method of claim 11 wherein said molecule
2 is DAB₄₈₆IL-2.

1 13. The method of claim 11 wherein said molecule
2 is DAB₃₈₉IL-4.


1 14. The method of claim 11 wherein said molecule
2 is DAB₃₈₉IL-6.

1 15. The method of claim 8 wherein said
2 interleukin is interleukin-4.

1 16. The method of claim 8 wherein said
2 interleukin is interleukin-6.

1

FIGURE 1.







FIGURE 3.



INTERNATIONAL SEARCH REPORT

International Application No. PCT/US92/01705

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ³		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC (5): A61K 37/02, 39/00, 39/12 US CL : 424/85.1, 85.2, 85.91, 89; 514/8; 435/69.52; 530/351		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁴		
Classification System	Classification Symbols	
U.S.	424/85.1, 85.2, 85.8, 85.91, 89; 514/2,8,21; 435/69.5,69.52,69.7,70.21; 935/106,107,109; 930/140,141; 530/351	
Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ⁵		
Computer data-base search of CAS and Dialog; For Treat/suppress/inhibit viral infection and Hybrid/Chimeric Protein Conjugate and Interleukin and toxin		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ¹⁴		
Category ⁶	Citation of Document, ¹⁸ with indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
X/Y	Cancer Research, Vol. 49, issued 15 July 1989, Kiyokawa et al., "Cytotoxicity in Interleukin 2-Toxin toward Lymphocytes From Patients with Adult T-Cell Leukemia," see pages 4042-46. See entire document.	1-12/1-12
X/Y	Nature, vol. 335, issued 22 September 1988, Chaudhary et al., "Selective Killing of HIV-infected Cells by Recombinant Human CD-4 Pseudomonas Exotoxin Hybrid Protein", page 369-372, see pages 369-72.	1-3, 6-7, 9-10/1-3, 6-7, 9-10
Y	WO, A, 91/101004, (Svrluga et al.), 24 January 1991, see entire document.	1-16
Y	Proc. Natl. Acad. Sci., vol. 86, issued June 1989, Ogata et al., "Cytotoxic Activity of a Recombinant Fusion Protein Between Interleukin-4 and Pseudomonas Exotoxin", see pages 4215-4219.	1-3, 6-10, 13, 15
Y	Proc. Natl. Acad. Sci., vol. 85, issued December 1988, Siegall et al., "Cytotoxic Activity of An Interleukin 6- Pseudomonas Exotoxin Fusion Protein on Human Myeloma Cells", see pages 9738-42.	1-3, 6-10, 14, 16
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>¹⁵ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search ²		Date of Mailing of the International Search Report ²
18 JUNE 1992		JUL 1992
International Searching Authority ¹		Signature of Authorized Officer ²⁰
ISA/US		GARNETTE D. DRAPER, PRIM. EXAM.

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y	European Journal Immunol., Vol. 20, issued April 1990, Waters et al., Interleukin-2 Receptor-Targeted Cytotoxicity. Receptor Binding Requirements for Entry of a Diphtheria Toxin-Related Interleukin 2 fusion Protein Into Cells," pages 785-91, see entire document	1-16
---	---	------

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

1. ☐ Claim numbers __, because they relate to subject matter (1) not required to be searched by this Authority, namely:
2. ☐ Claim numbers __, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out (1), specifically:
3. ☐ Claim numbers __, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING²

This International Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Search Authority did not invite payment of any additional fee.

Remark on protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.